Successful Treatment of Radiation-Induced Hemorrhagic Gastritis by Prednisolone

Feng-Woei Tsay, Gin-Ho Lo, Kwok-Hung Lai, Chiun-Ku Lin, Hoi-Hung Chen, and Ping-I Hsu

Division of Gastroenterology, Department of Internal Medicine, Kaohsiung Veterans General Hospita, and National Yang-Ming University, Kaohsiung, Taiwan, R.O.C

Abstract

Radiation-induced gastritis is a rare cause of gastrointestinal bleeding. Currently, the standard method for treatment of radiation-induced hemorrhagic gastritis has not been established. We presented a 53-year-old female patient with hemorrhagic gastritis and recurrent melena following external radiotherapy for cholangiocarcinoma. Endoscopic local injection and anti-secretary therapy were performed for hemostasis, but the symptoms still occurred off and on. Finally, high dose prednisolone was prescribed to her. The response was dramatic, and bleeding stopped 2 days later. She was then discharged with stable condition. There was no gastrointestinal bleeding in subsequent 5-month follow-up period. In conclusion, steroid therapy can effectively treat some patients with radiation-induced hemorrhagic gastroenteritis. (J Intern Med Taiwan 2006; 17: 39-42)

Key Words : Radiation, Hemorrhagic gastritis, Prednisolone

Introduction

Unlike radiation proctopathy, radiation-induced gastritis is a rare cause of gastrointestinal bleeding. We presented a 53-year-old female patient with hemorrhagic gastritis following external radiotherapy for cholangiocarcinoma. She was successfully treated by prednisolone therapy.

Case report

A 53-year-old woman was diagnosed as cholangiocarcinoma of right lobe liver with main portal vein thrombosis and lymphoadenopathy in March 1998. She received right lobectomy and lymphonode resection in April 1998. Four courses of intrahepatic arterial infusion chemotherapy with epirubicin, cisplatin and VP-16 were administered at our Department of General Surgery from April 1998 to September 1998. Besides she had also received several courses of chemotherapy and fractionated external beam radiotherapy at Sun Yet-Sen Cancer Center of Koo Foundation later. In December 1998, the patient began to suffer from intermittent melena, and she was admitted to our Gastrointestinal Ward two months later. On account of anemia (hemoglobin: 8.3 g/dL), blood transfusion was administered. A panendoscopy revealed diffuse hyperemic mucosa with telangiectasis from body of stomach to second portion of duodenum (figure 1). In addition, multiple hemorrhagic patches with active oozing were found over antrum (figure 2). Local injection and local spray with diluted epinephrine were done, but bleeding persisted. Lansoprazole and sucralfate were given in initial 1 month of hospital course. However, upper gastrointestinal bleeding with hypovolemic shock still occurred off and on. Under the impression of radiation-induced gastroenteritis, prednisolone, 40 mg daily, was prescribed to her, and its response was dramatic. Bleeding stopped 2 days later, and series blood examinations showed stable hemoglobin levels. She was discharged with stable condition 14 days following prednisolone therapy. Two months later she received follow-up panendoscopy, which revealed erythematous patches over antrum and body only. She received maintenance prednisone therapy (5 mg daily) later, and no gastrointestinal bleeding recurred in subsequent 5-months.

Discussion

Radiation therapy delivered to the upper abdomen may damage the gastric wall^{1,2}. The initial injury is characteristically acute inflammation of gastric mucosa. If injury progresses, vasculopathy of submucosa may occur and progress to obliterative endarteritis, vasculitis, and endothelial proliferation, leading to mucosal ischemia, ulceration, telangiectasia, and fibrosis ³⁻⁶.




Fig.1.Hyperemic mucosa with telangioectasis in the second portion of duodenum.




Fig.2.Multiple hemorrhagic patches with active oozing were found over antrum.

The diagnosis of radiation-induced gastritis is based on a high index of clinical suspicions for patient who received radiation therapy for upper abdomen lesions. The clinical presentations include epigastric pain, dysphagia, dyspepsia, heart burn sensation or melema. Endoscopy should be arranged for the patients presenting with aforementioned symptoms following radiation therapy for upper abdomi-

nal organs. The characteristic finding of radiation-induced gastric injury is the presence of telangiectasia, as seen in our patient. Other endoscopic findings include diffuse erythema of mucosa, shallow or deep ulcers and scar formation.

Currently, the golden standard method for treatment of radiation-induced hemorrhagic gastritis has not been established. Argon plasma coagulation is a new method of noncontact electrocoagulation in which high-frequency energy is delivered to the tissue through ionized argon. This technique is well suited for the coagulation of large bleeding surfaces and has the advantage of a limited depth of penetration (2 to 3 mm), thus minimizing risks for transmural necrosis and stricture formation, fistulization, and perforation⁷. It had been reported for successful hemostasis of radiation-induced hemorrhagic gastritis, colotis and proctitis 7,8,9. Epilon-aminocaproic acid, a fibrinolysis inhibitor which inhibits plasminogen activator, was reported effectively in control bleeding of radiation gastritis⁵. It helped fibrin deposits to control bleeding from multiple bleeding sites and mucosal ectasias in radiation-damaged gastrointestinal mucosa⁵. Kochhar et al. reported that steroids successfully treated radiation-induced proctosigmoiditis¹⁰, and Giuseppe et al. reported hydrocortisone prevented acute radiation proctitis during conformal radiotherapy for prostate carcinoma¹¹. However there is still insufficient data concerning the steroid effect on radiation-induced hemorrhagic gastritis.

The endoscopic injection therapy with diluted epinephrine plus anti-secretory therapy was initially done for hemostatsis in our patient with severe radiation gastroenteritis, but the effect was poor. At the time there was no machine of argon plasma coagulation in our hospital. We therefore tried high-dose prednisolone therapy, and it effectively controlled bleeding.

Radiation-induced severe hemorrhagic gastritis is rare. Currently argon plasma coagulation hemostasis is thought as more suited method. Due to the in-

accessibility of argon plasma coagulation, we therefore tried high-dose prednisolone therapy. It is effective in our patient. In conclusion, steroid therapy may effectively treat some patients with radiation-induced hemorrhagic gastroenteritis who have poor responses to endoscopic and anti-secretory therapies. However current data is still insufficient, and further large studies are needed.

References

- 1.Mascarenhas F, Silvestre ME, Sa da Costa M. Acute secondary effects in the esophagus in patients undergoing radiotherapy for carcinoma of lung. Am J Clin Oncol 1989; 12: 34-40.
- 2.Marrow JB, Dumot JA, Vargo JJ. Radiation-induced hemorrhagic carditis treated with argon plasma coagulator. Gastroinestest. Endosc 2000; 51: 498-9.
- 3.Earnest DL, Trier JS. Radiation enteritis and colitis. In: Sleiseng MH, Fordtran JS, eds. Gasteoinestinal Disease, 5th ed. Philadelphia: W.B. Saunders Co; 1993; 1257-68.
- 4.DeCosse JJ, Rhodes RS, Wents WB, Reagan JW, Dworken HJ, Holden WD. Natural history and management of radiation induced injury of the gastrointestinal tract. Ann Sur 1969; 170: 369.
- 5.Grover N, Jonhnson A. Aminocaproic acid used to control upper gastrointestinal bleeding in radiation gastritis. Digest Dis Sci 1997; 982-4.
- 6.Singh I, Laungani GB. Intravesical epeilon amminocaproic acid in management of intractable bladder hemorrhage. Urology 1992; 40: 227-9.
- 7.Silva RA, Correia AJ, Dias LM, Viana HL, Viana RL. Argon plasma coagulation therapy for hemorrhagic radiation proctosigmoiditis. Gastrointest Endosc 1999; 50: 221-4.
- 8.Wada S, Tamada K, Tomiyama T, Yamamoto H, Nakazawa K, Sugano K. Endoscopic hematostasis for radiation-induced gastritis using argon plasma coagulation. J Gastroentero Hepato 2003; 34: 1215-8.
- 9.Corbinais S, Garin L, Pagenault M, Bretagne JF. Successful treatment by argon plasma coagulation of bleeding radiation-induced gastroduodenal vasculopathy. Endoscopy 2002; 34: 593.
- 10. Kochhar R, Patel F, Dhar A, Sharma SC, Ayyagari S, Aggarwal R, Goenka MK, Gupta BD, Mehta SK. Radiation-induced proctosigmoiditis. Prospective, randomized, double-blind controlled trial of oral sulfasalazine plus rectal steroids versus rectal sucralfate. Digest Dis Sci 1991; 36: 103-7.
- 11.Sanguineti G, Franzone P, Marcenaro M, Foppiano F, Vitale V. Sucralfate versus mesalazine versus hydrocortisone in the prevention of acute radiation proctitis during conformal radiotherapy for prostate carcinoma. A randomized study. Strahlenther Onkol 2003; 179: 464-70.

以類固醇成功治癒放射線治療導致之 出血性胃炎:一病例報告

蔡峰偉 羅錦河 黎國洪 林俊谷 陳海雄 許秉毅 高雄榮民總醫院 内科部胃腸科 國立陽明大學

摘 要

臨床上因放射線治療導致的出血性胃炎並不常見,目前亦無有效的治療方式。本文報導一位53歲的膽管癌女性病患在接受放射線治療後,發生出血性腸胃炎,合併有反覆性的解黑便及出血性休克。在以內視鏡腎上腺素局部注射合併氫離子阻斷劑藥物治療下,仍有反覆出血的情形。在氫氣電漿凝固止血治療法無法獲得的情況下,最後我們嘗試著以高劑量口服類固醇施以治療。結果病患的病情迅速改善,於投藥第二天後便未再有出血情況,且於二週後順利出院。在爲期五個月的門診追蹤期間,並未再有出血情形。顯示類固醇對於控制某些放射治療後所引起的出血性胃腸炎有很好的效果。